

Critical species of Odonata in Australia

John H. Hawking¹ & Günther Theischinger²

¹ Cooperative Research Centre for Freshwater Ecology,
Murray-Darling Freshwater Research Centre, PO Box 921, Albury NSW,
Australia 2640. <john.hawking@csiro.au>

² Environment Protection Authority, New South Wales, 480 Weeroona Rd,
Lidcombe NSW, Australia 2141. <theischinger@epa.nsw.gov.au>

Key words: Odonata, dragonfly, IUCN, critical species, conservation, Australia.

ABSTRACT

The Australian Odonata fauna is reviewed. The state of the current taxonomy and ecology, studies on biodiversity, studies on larvae and the all identification keys are reported. The conservation status of the Australian odonates is evaluated and the endangered species identified. In addition the endemic species, species with unusual biology and species, not threatened yet, but maybe becoming critical in the future are discussed and listed.

INTRODUCTION

Australia has a diverse odonate fauna with many relict (most endemic) and most of the modern families (Watson et al. 1991). The Australian fauna is now largely described, but the lack of organised surveys resulted in limited distributional and ecological information. The conservation of Australian Odonata also received scant attention, except for Watson et al. (1991) promoting the awareness of Australia's large endemic fauna, the listing of four species as endangered (Moore 1997; IUCN 2003) and the suggesting of categories for all Australian species (Hawking 1999). This conservation report summarizes the odonate studies/literature for species found in Continental Australia (including nearby smaller and larger islands) plus Lord Howe Island and Norfolk Island. Australia encompasses tropical, temperate, arid, alpine and off shore island climatic regions, with the land mass situated between latitudes 11-44°S and 113-154°E, and flanked on the west by the Indian Ocean and on the east by the Pacific Ocean.

Table 1. Australian odonate species with unusual biology.

Family/species	Biology
Lestidae	
<i>Austrolestes minjerriba</i> Watson, 1979	Largely confined to brown acid coastal lakes
<i>Indolestes obiri</i> Watson, 1979	Breeds in rock pools in caves
<i>Lithosticta macra</i> Watson, 1991	Breeds in temporary streams draining rocky out crops
Megapodagrionidae	
<i>Podopteryx selysi</i> (Förster, 1899)	Larva lives in water-filled tree holes
Petaluridae	
<i>Petalura gigantea</i> Leach, 1815	Larva lives in water-filled tunnels, near water
<i>hesperia</i> Watson, 1958	Larva lives in water-filled tunnels, near water
<i>ingentissima</i> Tillyard, 1908	Larva lives in water-filled tunnels, near water
<i>pulcherrima</i> Tillyard, 1913	Larva lives in water-filled tunnels, near water
<i>litorea</i> Theischinger, 1999	Larva lives in water-filled tunnels, near water
Austropetaliidae	
<i>Austropetalia patricia</i> (Tillyard, 1910)	Larva lives on logs and moss or between rocks on the edges of waterfalls
<i>Austropetalia tonyana</i> Theischinger, 1995	Larva lives on logs and moss on the edges of waterfalls
Aeshnidae s.l.	
<i>Acanthaeschna victoria</i> Martin, 1901	Larva possibly inhabits coastal <i>Melaleuca</i> swamps
<i>Antipodophlebia asthenes</i> (Tillyard, 1916)	Last larval stage terrestrial
<i>Gynacantha nourlangie</i> Theischinger & Watson, 1991	Possibly breeds in rock pools in caves
<i>Telephlebia brevicauda</i> Tillyard, 1916	Larva semi-aquatic
<i>cyclops</i> Tillyard, 1916	Larva semi-aquatic
<i>godeffroyi</i> Selys, 1883	Larva semi-aquatic
<i>tillyardi</i> Campion, 1916	Larva semi-aquatic
<i>tryoni</i> Tillyard, 1917	Larva semi-aquatic
<i>undia</i> Theischinger, 1985	Larva semi-aquatic
Corduliidae s.l.	
<i>Pseudocordulia circularis</i> Tillyard, 1909	Larva lives in moist leaf litter
<i>elliptica</i> Tillyard, 1913	Larva lives in moist leaf litter
Libellulidae	
<i>Orthetrum boumiera</i> Watson & Arthington, 1978	Confined to brown acid coastal lakes

STATE OF THE ART

Studies on taxonomy and ecology

Starting with Fabricius (1775), many workers, most notably H. Burmeister, J.P. Rambur, F. Brauer, W.F. Kirby, E. de Selys Longchamps, Y. Sjöstedt, R.J. Tillyard, R. Martin, M.A. Lieftinck, F.C. Fraser, J.A.L. Watson, A.F. O'Farrell and G. Theischinger, accumulated taxonomic and some biological information on Australian Odonata, from which Houston & Watson (1988) compiled it in the Odonata section of the Zoological catalogue of Australia. Watson et al. (1991) included all the 301 known species in an illustrated book enabling the identification of the

adults and listed the major taxonomic revisions and studies on regional faunas, ecology and biodiversity of Australian Odonata. Watson & Houston (1994) published a checklist including the species and genera of Australian odonates and the relevant primary taxonomic literature.

Between 1991 and 2001, Carle (1995) and Theischinger (1998a, 1998b) established six genera, and Theischinger (1993, 1995a, 1995b, 1996a, 1996b, 1997a, 1997b, 1998b, 1998c, 1998d, 1998f, 1999a, 1999b, 1999c, 1999d, 2000c, 2001b, 2001c, 2003), Carle (1995) and Brown & Theischinger (1998) described 25 Australian species and subspecies. All these and other new taxonomic information were incorporated into a list of common (and scientific) names for all Australian odonate species (Hawking & Theischinger 2002) before the established records of *Nannophya pygmaea* Rambur, 1842 from Australia were referred to the newly established *N. paulsoni* (Theischinger 2003). Even more recently Theischinger (2004) attributed full generic status to two subgenera of *Austrogomphus* and, facing the uncertainty about *Austrogomphus praeruptus* and *A. melaleucae* Tillyard, 1909 being one or two species, stressed the existing urgency to get material of this complex/species from the far south of its distribution. Fresh larval evidence indicates the presence in Australia of *Selysioneura*, of an undescribed new species or genus close to *Austrophya mystica* and of a possibly undescribed species of *Nannophya* (GT unpubl.).

Detailed studies on the ecology of Odonata are scarce and the available information is generally natural history notes. Many of the early papers of R.J. Tillyard contained ecological information on the adults, larvae or both (Tillyard 1909a, 1909b, 1910a, 1910b, 1910c, 1911a, 1911b, 1912, 1913, 1916, 1928). Watson (1958, 1962, 1963, 1967, 1969) and Hodgkin & Watson (1958) presented ecology of Odonata from Western Australia, whereas Hawking (1986), Sant & New (1988), Hawking & Ingram (1994) and Hawking & New (1995a, 1995b, 1996, 1999, 2003) presented ecology of species from Victoria. Watson (1981) presented an overview of the ecology and biogeography of Australian Odonata, while Watson & Theischinger (1984) and Theischinger (2001, 2002) proposed and elaborated on, regions of taxonomic disjunctions.

Studies on biodiversity

Biodiversity studies of Australian Odonata are limited to particular areas or regions (Watson & Abbey 1980; Thompson 1989, 1991; Reeves & Woodall 1991; Wells & Cartwright 1992; Woodall 1992, 1993; Reeves 1993, 1995, 1998, 2003; Buck 1997, 1998; Davis 1997; Hawking 1998; Endersby 2000), islands (Reeves 1988a, 1988b, 1990; Endersby 2002), streams (Watson et al. 1978; Arthington & Watson 1982; Reeves 1987; O'Connor 1993; Hawking & New 1999, 2003) and unique habitats (Horwitz 1997). However, valuable information is available in the many major environmental surveys of invertebrates: Victoria (Smith et al. 1978; Malipatil & Blyth 1982; Marchant et al. 1984a, 1984b; Metzeling et al. 1984; Boulton & Lake 1992); New South Wales (Chessman & Williams 1999); Northern Territory (Marchant 1982; Davis 1997); Western Australia (Davis & Christidis 1997).

Identification guides

The Australian Odonata can be identified by Watson et al. (1991), however, this book does not include the 24 new taxa described since 1991 or specific information on the larvae. To address this we are currently preparing a new guide for the identification of adults and larvae of the Australian Odonata. Theischinger & Hawking (2003) produced an illustrated handbook for the identification of adults and larvae of Victoria and GT is also preparing an updated version of 'Tasmanian Odonata' (cf. Allbrook 1979).

Studies on larvae

In the past the poor state of larval taxonomy has been a major impediment to aquatic surveys, however, recent publications have started addressing this deficiency. Hawking (1993) complemented the biodiversity study of Northern Territory Odonata by Watson & Abbey (1980) based on adults only. Theischinger et al. (1993) studied the larvae of the Australian Synlestidae, Hawking & Theischinger (1999) of the entire order Odonata from New South Wales, and Theischinger (1998, 2000, 2001, 2002) of the Australian Gomphidae, Synthemiidae, Petaluridae, Archipetaliidae, Austropetaliidae, Aeshnidae s.l. and some Corduliidae s.l. For some of the more modern Corduliidae s.l. and the Libellulidae s.l. a larval identification and ecology guide is also being prepared by GT.

CRITICAL TAXA AND CONSERVATION CONCERNS

Notes on the species previously listed by IUCN

For Australia four species have been listed in the IUCN Red List of threatened species (IUCN 2003):

as critically endangered [CR]:

Austrocordulia leonardi

as endangered [EN]:

Petalura pulcherrima

as vulnerable [VU]:

Acanthaeschna victoria, *Hemiphlebia mirabilis*

Additionally, the following species were listed as "priority species" for Australia by Moore (1997):

as monotypic genera confined to one country only:

Chorismagrion risi, *Hemiphlebia mirabilis*, *Caliagrion billinghursti*, *Archipetalia auriculata*, *Austropetalia patricia*, *Acanthaeschna victoria*, *Antipodophlebia asthenes*, *Austrogynacantha heterogena*, *Austrophlebia costalis*, *Dendroaeschna conspersa*, *Armagomphus armiger*, 'Austroepigomphus' *praeruptus*, *Apocordulia macrops*, *Austrophya mystica*, *Hesperocordulia berthoudi*, *Pentathemis membranulata*, *Synthemiopsis gomphomacromioides*, *Austrothemis nigrescens*, *Notolibellula bicolor*.

Table 2. Odonate species, not threatened yet, but might become critical in the future, mainly taxonomically isolated species and monotypic genera endemic to Australia. TT: taxonomically isolated; ME: only species of monotypic, endemic genus.

Family/species	TT	ME	Threats and notes
Chorismagrionidae			
<i>Chorismagrion risi</i> Morton, 1914	○	●	Restricted range
Megapodagrionidae			
<i>Archargiolestes parvulus</i> (Watson, 1977)	○	○	Restricted range
<i>pusillissimus</i> Kennedy, 1925	○	○	Restricted range
<i>pusillus</i> (Tillyard, 1908)	○	○	Restricted range
<i>Miniargiolestes minimus</i> (Tillyard, 1908)	○	●	Restricted range
Isostictidae			
<i>Labidiosticta vallis</i> (Fraser, 1955)	○	●	Scattered distribution; habitat destruction
<i>Oristicta filicicola</i> Tillyard, 1913	○	●	
Coenagrionidae			
<i>Caligrion billinghursti</i> (Martin, 1901)	○	●	Scattered distribution; habitat destruction
<i>Coenagrion lyelli</i> (Tillyard, 1913)	●	○	
Diphlebiidae			
<i>Diphlebia coerulescens</i> Tillyard, 1913	○	○	Habitat destruction in south of range
<i>hybridoides</i> Tillyard, 1912	●	○	Restricted range
Lestodeidae			
<i>Lestoidea barbara</i> Watson, 1967	●	○	Restricted range
<i>lewisiana</i> Theischinger, 1996	○	○	Restricted range
Archipetaliidae			
<i>Archipetalia auriculata</i> Tillyard, 1917	○	●	Habitat destruction outside national parks
Austropetaliidae			
<i>Austropetalia patricia</i> (Tillyard, 1910)	○	○	Habitat destruction outside national parks
Aeshnidae s.l.			
<i>Antipodophlebia asthenes</i> (Tillyard, 1916)	○	●	Larva semi-terrestrial
<i>Austrogynacantha heterogena</i> Tillyard, 1908	○	●	
<i>Dendroaeschna conspersa</i> (Tillyard, 1907)	○	●	
Gomphidae			
<i>Austrogomphus lateralis</i> (Selys, 1873)	●	○	
Synthemistidae			
<i>Austrosynthemis cyanitincta</i> (Tillyard, 1908)	○	●	
<i>Parasynthemis regina</i> (Selys, 1874)	○	●	
<i>Synthemopsis gomphomacromioides</i> Tillyard, 1917	○	●	
Corduliidae s.l.			
<i>Apocordulia macrops</i> Watson, 1980	○	●	Habitat destruction
<i>Cordulephya montana</i> Tillyard, 1911	●	○	Distribution scattered
<i>Pentathemis membranulata</i> Karsch, 1890	○	●	Scattered distribution
Libellulidae			
<i>Austrothemis nigrescens</i> (Martin, 1901)	○	●	Habitat destruction
<i>Notolibellula bicolor</i> Theischinger & Watson, 1977	○	●	Scattered distribution

as taxonomically isolated species:

Diphlebia coerulescens, *D. euphoeoides* Tillyard, 1907, *D. hybridoides*, *D. lestoides* (Selys, 1853), *D. nymphoides* Tillyard, 1912, *Chorismagrion risi*, *Lestoidea barbarae* Watson, 1967, *L. conjuncta* Tillyard 1913, *Hemiphlebia mirabilis*, *Petalura gigantea*, *P. hesperia*, *P. ingentissima*, *P. pulcherrima*, *Archipetalia auriculata*, *Austropetalia patricia*, *Cordulephya bidens* Sjöstedt, 1917, *C. divergens*, *C. montana*, *C. pygmaea*.

The species listed by the IUCN (2003) are commented on in the next chapter. Concerning the species listed by Moore (1997) it has to be noted that the genus *Austroepigomphus* is synonym to *Austrogomphus*.

Table 3. Suggested Red List categories (SRL) for critical species. CR: critically endangered; EN: endangered; VU: vulnerable; NT: near threatened.

Family/species	SRL	Threats and notes
Megapodagrionidae		
<i>Griseargiolestes bucki</i> Theischinger, 1998	VU	Restricted range
Hemiphlebidae		
<i>Hemiphlebia mirabilis</i> Selys, 1869	VU	Habitat destruction
Isostictidae		
<i>Lithosticta macra</i> Watson, 1991	VU	Restricted range
Petaluridae		
<i>Petalura gigantea</i> Leach, 1815	VU	Habitat destruction
<i>hesperia</i> Watson, 1958	VU	Habitat destruction
<i>ingentissima</i> Tillyard, 1908	VU	Habitat destruction
<i>litorea</i> Theischinger, 1999	VU	Habitat destruction
<i>pulcherrima</i> Tillyard, 1913	EN	Status uncertain
Aeshnidae		
<i>Acanthaeschna victoria</i> Martin, 1901	EN	Habitat destruction
Gomphidae		
<i>Antipodogomphus dentosus</i> Watson, 1991	EN	Restricted range
<i>hodgkini</i> Watson, 1969	EN	Restricted range
<i>Armogomphus armiger</i> (Tillyard, 1913)	VU	
<i>Austrogomphus gordonii</i> Watson, 1962	VU	Restricted range
<i>longipositor</i> (Watson, 1991)	VU	Scattered distribution
<i>praeruptus</i> (Selys, 1857)	EN	Known only from holotype
<i>Hemigomphus atratus</i> Watson, 1991	VU	Known only from holotype
<i>magela</i> Watson, 1991	VU	Restricted Range
<i>Odontogomphus donnellyi</i> Watson, 1991	VU	Restricted range
Synthemistidae		
<i>Archaeosynthemis spiniger</i> (Tillyard, 1913)	VU	Very few specimens known

Family/species	SRL	Threats and notes
Corduliidae s.l.		
<i>Archaeophya adamsi</i> Fraser, 1959	CR	Habitat destruction
<i>Austrocordulia leonardi</i> Theischinger, 1973	CR	Habitat destruction, disappeared from several localities
<i>Cordulephya divergens</i> Tillyard, 1917	VU	Known only from the Sydney area
<i>Hesperocordulia berthoudi</i> Tillyard, 1911	VU	
<i>Lathrocordulia garrisoni</i> Theischinger & Watson, 1991	VU	Known only from holotype
Libellulidae		
<i>Huonia melvillensis</i> Brown & Theischinger, 1998	VU	Known only from a single locality

Species to be considered

Hemiphysalia mirabilis

It is probably the world's most primitive odonate, retaining venation characteristic of the Permian Period. *H. mirabilis* was thought to be extinct until rediscovered by Davies (1985) at Wilsons Promontory, Victoria. Later it was re-discovered at its original location at Alexandra, Victoria and it is now known from six small, scattered populations ranging from central eastern Victoria to Northern Tasmania (Trueman et al. 1992; Endersby 1993a, 1993b). Its biology and conservation were studied by Sant & New (1988) and by New (1993), its habitat is threatened by grazing and burning (Watson 1995). This species is still listed as 'endangered' and is protected under the Victorian Government Flora and Fauna Guarantee Act 1988 (Birkin & Quin 1993).

Genus *Petalura*

Members of the ancient family Petaluridae were abundant in the Jurassic, 190 million years ago (Carpenter 1992) and possibly five of the 11 extant species occur in Australia. They all belong in the endemic genus *Petalura* and include some of the largest and most spectacular odonates of the world. They are of conservation concern due to the unique life-style of the larvae, being semi-aquatic, having a long life-cycle and inhabiting permanent burrows excavated in swampy ground. Presently five species are known, *P. hesperia* from south-western Australia, *P. gigantea* from New South Wales, *P. litorea* from south-eastern Queensland and north-eastern New South Wales, and *P. ingentissima* and *P. pulcherrima* from north-eastern Queensland. The IUCN (2003) red-listed *P. pulcherrima* as 'endangered', however, its status is uncertain, and Davies (1998) suggested it was "awaiting rediscovery and confirmation of specific status".

In 2000-2001 National Parks & Wildlife Service funded searches for *P. gigantea* in eastern New South Wales, and these confirmed the distribution of *P. gigantea* over a large geographical range, but in a small number of sites and usually in small populations (J. Trueman pers. comm.; GT unpubl.). Subsequently, *P. gigantea* was finally listed as an 'endangered' species under Schedule 1, Part 1, of the 'Threatened Species Conservation Act' in New South Wales; <<http://www.national-parks.nsw.gov.au/npws.nsf/Content/Giant+dragonfly+-+endangered+species+listing>>.

P. gigantea is reasonably secure occurring in areas now protected as national parks and reserves. *P. ingentissima* is wide-spread, occurring in the Wet Tropics World Heritage zone of northern Queensland and is protected in the Heritage area. However, *P. litorea* is known from a small number of sites in southern Queensland and from only two in New South Wales. The aquifer under Stradbroke Island where the largest and best-known population still exists is being “pumped” to provide water for Brisbane. *P. hesperia* was known from nine populations, but is currently recorded from only six populations in south-western Australia (Barrett 1996). The species’ patchy distribution subjects it to threats from land clearing for agriculture and bauxite mining.

Acanthaeschna victoria

Until fairly recently *A. victoria* was not collected or seen alive by collectors of odonates, and only a few specimens were collected, these by non-specialists, over more than 100 years. In 1999, however, several individuals were collected/observed in coastal swampland near Broadwater in north-eastern New South Wales. Several weeks later this habitat was found completely destroyed by bulldozers (Theischinger 2000b). However, since then single individuals have been observed in three more coastal swamps, situated at some distance from each other, in north-eastern New South Wales. Draining of the coastal fringe for housing, agriculture and entertainment still remain the biggest threat to *A. victoria*, especially as its behaviour and ecology are still unknown. Habitat reduction and fragmentation have probably threatened this species since its original discovery in the 19th century.

Archaeophya adamsi

Only the type locality in south-eastern Queensland and very few localities, all within 100 km of Sydney, were previously known for *A. adamsi* (Theischinger & Watson 1977, 1982; Hawking & Theischinger 1999). It was gazetted on 12 November 1999 as a vulnerable species in New South Wales and now legally protected (NSW Fisheries 2002) (<http://www.fisheries.nsw.gov.au/thr/species/fn-adams-emerald-dragonfly.htm>). However, in 2002 two adults were collected in the foothills of the Blue Mountains (GT unpubl.) and recently larvae were found in a few creeks near Jamison, New South Wales. Larvae were also collected in 2000 in Tunks Creek near Galston Gorge where the species was collected in 1969 and was supposed to have disappeared since. Fewer than ten adults and only a few larvae have ever been collected, even with extensive searches. This inability to find specimens, the occurrence of only six localised populations and major threats from rapid housing development from the expansion of Sydney have resulted in action to protect *A. adamsi*. It is now listed as a ‘vulnerable’ species in New South Wales and legally protected.

Austrocordulia leonardi

It was discovered in 1968 from Woronora River and Kangaroo Creek, south of Sydney (Theischinger 1973) and later recorded from the Nepean River, Maldon

Bridge near Wilton (Theischinger 1997c; Hawking & Theischinger 1999). This species has always been rare (only 11 adult specimens are known) and its occurrence is generally only recorded from the presence of exuviae. However, intensive surveys by GT over the last two years have failed to detect the presence of any of the life stages along Woronora River and Kangaroo Creek. *A. leonardi* was red-listed by Moore (1997) as 'critically endangered' due to its rareness and restricted nature. This listing is extremely relevant with the disappearance of the species from the type locality, along the Woronora River, after the removal of the weir at Heathcote. All specimens ever collected come from deep and shady man-made riverine pools with cooler water (along Woronora River, Kangaroo Ck and Nepean River). Natural deep pools probably disappeared some time ago with the creation of large dams.

Endemism

Compared to its size, Australia's odonate fauna has a very high share of endemism at familial, generic and specific level. According to Watson et al. (1991) almost 50% of the species appear to be Gondwanic in origin and of the 325 now recognised species of Australian Odonata, some 245 or 75% are endemic, and most of these are distributed over relatively small areas, or even site restricted.

Table 4. Critical Odonata found in the Australian region, also data deficient species, which might have to be deleted from the list. DD: data deficient; RR: range restricted; IC: identity of species needs clarification; A: action recommended.

Family/species	DD	RR	IC	A	Distribution and notes
Calopterygidae					
<i>Neurobasis australis</i> Selys, 1897	●	?	○	○	Status of (single) record uncertain
Chlorocyphidae					
<i>Rhinocypha tinca semitincta</i> Selys, 1869	●	?	○	○	Only records from 19th century
Synlestidae					
<i>Episynlestes intermedius</i> Theischinger & Watson, 1985	●	●	○	○	Possibly confined to Eungella area, Qld
Lestidae					
<i>Austrolestes aleison</i> Watson & Moulds, 1979+	●	○	○	○	
<i>Indolestes alleni</i> (Tillyard, 1913)	●	○	○	○	
Lestoideidae					
<i>Lestoidea barbara</i> Watson, 1967	●	●	○	○	Known from a few specimens
<i>lewisiana</i> Theischinger, 1996	●	●	○	○	Possibly restricted to Mt Lewis, Qld
Megapodagrionidae					
<i>Griseargiolestes bucki</i> Theischinger, 1998	●	○	○	○	Scattered distribution in ne NSW
<i>fontanus</i> (Tillyard, 1913)	●	○	○	○	Scattered distribution. Known from only a few localities
<i>metallicus</i> (Sjöstedt, 1917)	●	○	○	○	Confined to tropical rainforests in Qld
<i>Austroargiolestes alpinus</i> (Tillyard, 1913)	●	○	○	○	Scattered distribution
<i>brookhousei</i> Theischinger & O'Farrell, 1986	●	○	○	○	Scattered distribution
<i>christine</i> Theischinger & O'Farrell, 1986	●	○	○	○	Scattered distribution
<i>chrysoides</i> (Tillyard, 1913)	●	○	○	○	
<i>elke</i> Theischinger & O'Farrell, 1986	●	●	○	○	Possibly confined to Eungella area, Qld

Family/species	DD	RR	IC	A	Distribution and notes
Hemiphlebiidae					
<i>Hemiphebia mirabilis</i> Selys, 1869	○	●	○	●	Listed as "endangered" in Victoria; IUCN Red List (VU)
Coenagrionidae					
<i>Agriocnemis dobsoni</i> Fraser, 1954	●	○	○	○	
<i>rubricauda</i> Tillyard, 1913	●	○	○	○	
<i>thoracalis</i> Sjöstedt, 1917	●	○	●	○	Single male known
<i>Austrocnemis obscura</i> Theischinger & Watson, 1901	●	●	○	○	
Isostictidae					
<i>Austrosticta fieldi</i> Tillyard, 1908	●	○	○	○	
<i>frater</i> Theischinger, 1997	●	●	○	○	Known only from two specimens
<i>soror</i> Sjöstedt, 1917	●	○	○	○	
<i>Eurysticta kununurra</i> Watson, 1991	●	○	○	○	Known from only a few localities
<i>Lithosticta macra</i> Watson, 1991	○	●	○	●	Noted as critical in environmental reports of ERISS ¹
<i>Neosticta silvarum</i> (Sjöstedt, 1917)	●	○	○	○	Scattered distribution
Protoneuridae					
<i>Nososticta kalumburu</i> Watson & Theischinger, 1984	○	●	○	○	Confined to the Kimberley, W.A.
<i>koolpinyah</i> Watson & Theischinger, 1984	?	?	○	○	Known from only a few localities
Petaluridae					
<i>Petalura gigantea</i> Leach, 1815	●	○	○	●	Listed as "endangered" in NSW
<i>hesperia</i> Watson, 1958	○	●	○	●	Surveys have been undertaken
<i>ingentissima</i> Tillyard, 1908	●	○	○	○	Identification of some material doubtful because of status problem in <i>P. pulcherrima</i>
<i>litorea</i> Theischinger, 1999	●	○	○	○	Widely scattered distribution
<i>pulcherrima</i> Tillyard, 1913	●	?	●	●	Status uncertain; IUCN Red List (EN)
Austropetaliiidae					
<i>Austropetalia tonyana</i> Theischinger, 1995	●	○	○	○	Restricted distribution
Aeshnidae s.l.					
<i>Acanthaeschna victoria</i> Martin, 1901	●	○	○	●	Apparently scattered distribution due to habitat destruction; IUCN Red List (VU)
<i>Austroaeschna christine</i> Theischinger, 1993	●	●	○	○	Possibly confined to Eungella area, Qld; threatened by dairy farming
<i>eungella</i> Theischinger, 1993	●	●	○	○	Possibly confined to wider Eungella area, Qld; threatened by dairy farming
<i>flavomaculata</i> Tillyard, 1916	○	●	○	○	Restricted to the snow areas of southern alps
<i>muelleri</i> Theischinger, 1982	●	●	○	○	Possibly restricted to Carnarvon Gorge, Qld
<i>Austrophlebia subcostalis</i> Theischinger, 1996	●	○	○	○	Scattered distribution in tropical Qld
<i>Anax georgius</i> Selys, 1872	●	●	○	○	Possibly restricted to the Kimberley
<i>Spinaeschna watsoni</i> Theischinger, 1982	●	○	○	○	Only two adults available
<i>Telephlebia cyclops</i> Tillyard, 1916	●	○	○	○	Confined to rainforest in NSW and Qld
<i>tillyardi</i> Campion, 1916	●	○	○	○	Confined to rainforests in tropical Qld
<i>tryoni</i> Tillyard, 1917	●	○	○	○	Possibly confined to coastal situations in NSW and southern Qld
<i>undia</i> Theischinger, 1985	●	●	○	○	Possibly restricted to Carnarvon Gorge, Qld

Family/species	DD	RR	IC	A	Distribution and notes
Gomphidae s.l.					
<i>Antipodogomphus dentosus</i> Watson, 1991	○	●	○	○	Possibly restricted to South Alligator and Katherine Rivers
<i>edentulus</i> Watson, 1991	●	●	○	○	Only a few females known
<i>hodgkini</i> Watson, 1969	●	●	○	○	Restricted to nw Australia
<i>Armagomphus armiger</i> (Tillyard, 1913)	○	●	○	○	Restricted distribution in sw Australia
<i>Austrogomphus angelorum</i> Tillyard, 1913	●	○	○	○	Known from fewer than 10 specimens; only found along the Murray River, larva unknown
<i>bifurcatus</i> Tillyard, 1909	●	●	○	○	
<i>divaricatus</i> Watson, 1991	●	●	○	○	
<i>doddi</i> Tillyard, 1909	●	○	○	○	
<i>gordoni</i> Watson, 1962	○	●	○	○	Restricted distribution
<i>longipositor</i> (Watson, 1991)	●	○	○	○	Known from a few females
<i>mouldsorum</i> Theischinger, 1999	●	●	○	○	Probably confined to the Kimberley; Known only from a single female
<i>praeruptus</i> (Selys, 1857)	●	●	●	○	Known from the holotype. Possibly the same species as <i>A. melaleuca</i> Tillyard, 1909
<i>pusillus</i> Sjöstedt, 1917	●	○	○	○	Single male known
<i>Hemigomphus atratus</i> Watson, 1991	●	○	○	○	Known only from the holotype
<i>cooloola</i> Watson, 1991	●	●	○	○	Apparently confined to Cooloola area and Fraser and Stradbroke Islands
<i>magela</i> Watson, 1991	●	●	○	○	Restricted distribution
<i>theischingeri</i> Watson, 1991	●	●	○	○	Confined to rainforests in tropical Qld
<i>Ictinogomphus paulini</i> Watson, 1991	●	●	○	○	Only a few males are known
<i>Odontogomphus donnellyi</i> Watson, 1991	●	●	○	○	Apparently confined to a small area in tropical Qld
Synthemistidae					
<i>Archaeosynthemis spiniger</i> (Tillyard, 1913)	●	●	○	○	Very few specimens known
<i>Choristhemis olivei</i> (Tillyard, 1909)	●	●	○	○	Very few specimens known
<i>Eusynthemis barbarae</i> (Moulds, 1985)	●	●	○	○	Apparently confined to Mt Lewis area, Qld; very few specimens known
<i>deniseae</i> Theischinger, 1977	●	●	○	○	Apparently confined to the Carnarvon Range, Qld
<i>netta</i> Theischinger, 1999	●	●	○	○	Apparently confined to Mt Lewis area, Qld; only three specimens known
<i>rentziana</i> Theischinger, 1998	●	○	○	○	?Scattered distribution
<i>tenera</i> Theischinger, 1995	●	●	○	○	Very few specimens known
<i>ursa</i> Theischinger, 1999	●	●	○	○	Possibly confined to high altitude on Barington Tops, NSW
<i>ursula</i> Theischinger, 1998	●	●	○	○	Possibly confined to Nothofagus forest in Chichester State Forest, NSW
<i>Tonyosynthemis claviculata</i> (Tillyard, 1909)	●	○	○	○	
<i>ofarrelli</i> (Theischinger & Watson, 1986)	●	○	○	○	Known only from a few specimens
Corduliidae s.l.					
<i>Archaeophya adamsi</i> Fraser, 1959	●	●	○	●	Listed as "vulnerable" in NSW
<i>magnifica</i> Theischinger & Watson, 1978	●	○	○	○	Systematically significant species,
<i>Austrocordulia leonardi</i> Theischinger, 1973	○	●	○	○	IUCN Red List (CE)

Family/species	DD	RR	IC	A	Distribution and notes
Corduliidae s.l. (continued)					
<i>Austrophya mystica</i> Tillyard, 1909	●	○	○	○	Confined to tropical rainforests in Qld
<i>Cordulephya divergens</i> Tillyard, 1917	○	●	○	○	Known only from the Sydney area
<i>Hemicordulia kalliste</i> Theischinger & Watson, 1991	●	●	○	○	
<i>novahollandiae</i> (Selys, 1871)	●	?	●	○	Taxonomic interpretation not possible at the present
<i>Hesperocordulia berthoudi</i> Tillyard, 1911	●	○	○	○	
<i>Lathrocordulia garrisoni</i> Theischinger & Watson, 1991	●	●	○	○	Known only from the holotype
<i>metallica</i> Tillyard, 1911	●	○	○	○	
<i>Macromia viridescens</i> Tillyard, 1911	●	●	○	○	Confined to Iron Range
<i>Metaphya tillyardi</i> Ris, 1913	●	●	○	○	Only a single specimen from an Australian island off the Papuan coast
<i>Micromidia rodericki</i> Fraser, 1959	●	●	○	○	Possibly extinct (habitat destruction)
Libellulidae					
<i>Austrothemis nigrescens</i> (Martin, 1901)	○	●	○	○	Monotypic genus; ?confined to coastal acid lakes;
<i>Camacinia othello</i> Tillyard, 1908	●	●	○	○	
<i>Huonia melvillensis</i> Brown & Theischinger, 1998	●	●	○	○	Known only from type locality on Melville Is.
<i>Nannophlebia mudginberri</i> Watson & Theischinger, 1991	○	●	○	○	Confined to Kakadu National Park
<i>Orthetrum boumiera</i> Watson & Arthington, 1978	○	●	○	○	Confined to brown acid coastal lakes; Threat from housing development
<i>serapia</i> Watson, 1984	●	○	○	○	
<i>Tetrathemis irregularis cladophila</i> Tillyard, 1908	●	●	○	○	Confined to tropical rainforests in Qld
<i>Trapezostigma¹ propinqua</i> (Lieftinck, 1942)	●	○	○	○	Habitats unknown

¹ Environmental Research Institute of Supervising Scientist, NT ² Syn. *Tramea*

Seven taxa treated variously as families or subfamilies in recent systematic papers (Watson et al. 1991; Carle 1995; Bechly 1996; Lohmann 1996) are endemic for Australia: Chorismagrionidae, Lestoideidae, Hemiphlebiidae, Archipetaliidae, Austropetaliidae, Pseudocorduliidae, Cordulephyinae. All are also monotypic at the generic level, and three of them even at the specific level. Independent of their current or future recognition at this or any other taxonomic level, these groups are significant for the understanding of evolution and relationships within the Odonata. In the currently recognised 110 genera of Australian Odonata, 49 are endemic, 22 of which are also monotypic. Monotypic endemic Australian genera comprise: *Miniargiolestes*, *Chorismagrion*, *Hemiphlebia*, *Labidiosticta*, *Lithosticta*, *Oristicta*, *Acanthaeschna*, *Antipodophlebia*, *Archipetalia*, *Dendroaeschna*, *Armagomphus*, *Odontogomphus*, *Apocordulia*, *Austrophya*, *Austrosynthemis*, *Hesperocordulia*, *Parasynthemis*, *Pentathemis*, *Synthemioptis*, *Austrothemis*, and *Notolibellula*. Genera are increasingly recognised as the most useful supra-specific taxon on which to base phylogenetic studies and these endemic and monotypic taxa have high predictive value for environmental assessment studies.

Species with unusual biology

Australia has a wide range of aquatic and semi-aquatic habitats; therefore it is not surprising that 23 species are different to most Odonata because of their unusual biology and unique habitats (Table 1). They range from species like *Antipodophlebia asthenes* whose final larval stage lives in rainforest leaf-litter (Watson & Theischinger 1980), to species of *Telephlebia* which have semi-aquatic larvae and *Indolestes obiri* which lives in pools in caves (Watson et al. 1991).

Species, not threatened yet, but might become critical in the future

Twenty-seven species qualify for near threatened (Table 2), as they could become critical if endangered in the future due to environmental degradation. Sixteen species are monotypic, endemic genera. Five species are taxonomically isolated. Most species have restricted ranges or scattered distributions and are confined to streams or bogs.

Suggested Red List category

Twenty-five species are deemed critical (Table 3). Two species should be listed as critically endangered, five as endangered and 18 as vulnerable.

General on conservation

Most of the knowledge about distributions and habitats of Australian Odonata has been gathered by naturalists, in their own time, and is largely anecdotal, but has provided the basis of most of Australia's Odonata knowledge. The lack of properly designed, commissioned surveys by professional researchers has resulted in the high proportion of 'data deficient' species, approximately 30% (Table 4), and is possibly the major concern facing the Australian Odonata conservation. Another concern is the export of the rare, unprotected fauna. According to D.A.L. Davies (in litt.), there is considerably more material of tropical *Petalura* in overseas collections than in Australian collections. It appears that the export control for *Petalura* was and is not effective and should be stepped up, because recently the purchase price of *P. ingentissima* on the "Japanese market" dropped manyfold (H. Karube pers. comm.).

Hawking (1999) presented a comprehensive paper on conservation issues for Australian Odonata and evaluated the conservation status for all known species and suggested IUCN categories for these species. Since then, two species, *Archaeophya adamsi*, and *P. gigantea* have been listed as vulnerable/endangered in New South Wales, and *Hemphlebia mirabilis* has been listed as endangered in Victoria. In 2000-2001 National Parks & Wildlife Service funded searches for *P. gigantea* in eastern New South Wales, and a colour brochure on the "Giant dragonfly" was handed out to the public.

CURRENT ACTIVITIES

Between 1994 and 2000 a nationwide program MRHI (Monitoring River Health Initiative) was conducted Australia-wide. During this study which amassed a large quantity of macroinvertebrate material from several thousand sites, many odonate larvae were collected, preserved and identified to familial level. There is now a funded project to identify at least some of them to species and to establish ecological species profiles. Similar studies conducted by the EPA (Environment Protection Authority) New South Wales on the Sydney Catchment, by the EPA Victoria on Victorian streams and by the Australian Water Quality Centre on South Australian streams have identified the odonate larvae to specific level.

The data basing of adult Odonata in the Australian National Insect Collection - Australia's largest Odonata collection - is currently processing the final group, the libellulids. This data base will be a valuable resource to odontologists, however, presently there are no plans to data base the odonates in the collections of other museums.

Colour photographs and conservation information on *Petalura gigantea*, *Apocordulia macrops* and *Archaeophya adamsi* have been published in newspaper articles and brochures handed out to the public on water quality and conservation issues. The New South Wales government is trying to raise public awareness of the conservation of odonates and promote them as indicators of a healthy environment.

The Australian Dragonfly Society was formed in February 2000 in order to disseminate information on Odonata of Australia and south-west Pacific. The society produces a newsletter 'Austrolestes' several times a year and it is an important organ to disseminate technical information, raise conservation concerns and provide naturalist notes on distribution. The Worldwide Dragonfly Association "Symposium of Odonatology" was held at Beechworth, Victoria (January 2003) and the post-symposium collecting trip supplied a good species list of several areas in Victoria and provided valuable information regarding the conservation status of *Telephlebia brevicauda*.

REFERENCES

- Allbrook, P., 1979. Tasmanian Odonata. University of Tasmania, Hobart.
- Arthington, A.H. & J.A.L. Watson, 1982. Dragonflies (Odonata) of coastal sand-dune fresh waters of south-eastern Queensland and north-eastern New South Wales. Australian Journal of Marine and Freshwater Research 33: 77-88.
- Barrett, M.D., 1996. Identification of significant habitats occupied by the Western *Petalura* dragonfly, *Petalura hesperia*, in the south-west of Western Australia. Unpublished report to the Australian Heritage Commission, Perth.
- Bechly, G., 1996. Morphologische Untersuchungen am Flügelgeäder der rezenten Libellen und deren Stammgruppenvertreter (Insecta; Pterygota; Odonata) unter besonderer Berücksichtigung der Phylogenetischen Systematik und des Grundplanes der Odonata. *Petalura*, Special Volume 2: 1-402.

- Birkin, E. & B. Quin, 1993. *Hemiphlebia* Damselfly (*Hemiphlebia mirabilis*) (Flora and Fauna Guarantee Action Statement No. 46). Department of Conservation and Environment, Melbourne.
- Boulton, A.J. & P.S. Lake, 1992. The macroinvertebrate assemblages in pools and riffles in two intermittent streams (Werribee and Lerderderg rivers, southern central Victoria). Occasional Papers of the Museum of Victoria 5: 55-71.
- Brown, G.R. & G. Theischinger, 1998. A new species of *Huonia* Förster from Australia (Anisoptera: Libellulidae). Odonatologica 27: 99-103.
- Buck, K., 1997. Drei Reisen nach New South Wales. Hagenia 14: 23-27.
- Buck, K., 1998. Odonatological notes from three trips to New South Wales, Australia. Notulae Odonatologicae 5: 7-8.
- Carle, F.L., 1995. Evolution, taxonomy, and biogeography of ancient Gondwanian Libelluloidea, with comments on anisopteroid evolution and phylogenetic systematics (Anisoptera: Libelluloidea). Odonatologica 24: 383-424.
- Carpenter, F.M., 1992. Treatise on invertebrate paleontology, Part R, Arthropoda 4, volume 3: superclass Hexapoda. The Geological Society of America, Inc. & Boulder, Colorado, and Lawrence, Kansas.
- Chessman, B.C. & S.A. Williams, 1999. Biodiversity and conservation of river macroinvertebrates on an expanding urban fringe: western Sydney, New South Wales, Australia. Pacific Conservation Biology 5: 36-55.
- Davies, D.A.L., 1985. *Hemiphlebia mirabilis* Selys: some notes on distribution and conservation status (Zygoptera: Hemiphlebiidae). Odonatologica 14: 331-339.
- Davies, D.A.L., 1998. The genus *Petalura*: field observations, habitats and conservation status (Anisoptera: Petaluridae). Odonatologica 27: 287-305.
- Davis, J., 1997. Conservation of aquatic invertebrate communities in central Australia. Memoirs of the Museum of Victoria 56: 491-503.
- Davis, J. & F. Christidis, 1997. A guide to wetland invertebrates of south-western Australia. Western Australian Museum, Perth.
- Endersby, I.D., 1993a. A new locality record for *Hemiphlebia mirabilis* Selys (Zygoptera: Hemiphlebiidae). Notulae Odonatologicae 4: 31-32.
- Endersby, I.D., 1993b. A new locality for *Hemiphlebia mirabilis* Selys (Odonata: Hemiphlebiidae). Victorian Entomologist 23: 4-5.
- Endersby, I.D., 2000. Checklist of Victorian dragonflies (Insecta: Odonata). Proceedings of the Royal Society of Victoria 112: 59-64.
- Endersby, I.D., 2002. The dragonflies of Norfolk Island, with the first record of *Pantala flavescens* (F.) (Odonata: Libellulidae). Entomologist's Monthly Magazine 138: 241-246.
- Fabricius, J.C., 1775. V. Vnogata. In: "Systema entomologiae, sistens insectorum classes, ordines, genera, species, adiectis synonymis, locis, descriptionibus, observationibus", Kortius, Flensburg & Leipzig, pp. 420-426.
- Hawking, J.H., 1993. A preliminary guide to the identification of dragonfly larvae (Odonata) from the Alligator Rivers Region of the Northern Territory. Open File Record 102. Office of the Supervising Scientist, Alligator Rivers Region, Jabiru.
- Hawking, J.H., 1998. The conservation status of dragonflies (Odonata) from south-eastern Australia. Memoirs of the Museum of Victoria 56: 537-542.

- Hawking, J.H., 1999. An evaluation of the current conservation status of Australian dragonflies (Odonata). In: Ponder, W. & D. Lunney (eds) "The other 99%. The conservation and biodiversity of invertebrates", Transactions of the Royal Zoological Society of New South Wales, Mosman, pp. 354-360.
- Hawking, J.H., 1998. Freshwater macroinvertebrates of Mount Buffalo. The Victorian Naturalist 115: 215-221.
- Hawking, J.H. & B.A. Ingram, 1994. Rate of larval development of *Pantala flavescens* (Fabricius) (Odonata: Libellulidae) at its southern limit of range. Odonatologica 23: 63-68.
- Hawking, J.H. & T.R. New, 1995a. The diet of anisopteran larvae from two streams in north-eastern Victoria, Australia. Odonatologica 24: 115-122.
- Hawking, J.H. & T.R. New, 1995b. The development of eggs of dragonflies (Odonata: Anisoptera) from two streams in north-eastern Victoria, Australia. Aquatic Insects 17: 175-180.
- Hawking, J.H. & T.R. New, 1996. The development of dragonfly larvae (Odonata: Anisoptera) from two streams in north-eastern Victoria, Australia. Hydrobiologia 317: 13-30.
- Hawking, J.H. & T.R. New, 1999. The distribution patterns of dragonflies (Insecta: Odonata) along the Kiewa River, Australia, and their relevance in conservation assessment. Hydrobiologia 392: 249-260.
- Hawking, J.H. & T.R. New, 2003. Interpreting dragonfly diversity to aid in conservation assessment: lessons from the Odonata assemblage at Middle Creek, north-eastern Victoria, Australia. Journal of Insect Conservation 6: 171-178.
- Hawking, J.H. & G. Theischinger, 1999. Dragonfly larvae (Odonata). A guide to the identification of larvae of Australian families and to the identification and ecology of larvae from New South Wales. Cooperative Research Centre for Freshwater Ecology, Thurgooda & Australian Water Technologies Pty Ltd, West Ryde.
- Hawking, J.H. & G. Theischinger, 2002. Vernacular names for the Australian dragonflies (Odonata). Austrolestes, Supplement to Number 4: 1-6.
- Hodgkin, E. P. & J.A.L. Watson, 1958. Breeding of dragonflies in temporary waters. Nature, London 181: 1015-1016.
- Horwitz, P., 1997. Comparative endemism and richness of the aquatic invertebrate fauna in peatlands and shrublands of far south-western Australia. Memoirs of the Museum of Victoria 56: 313-321.
- Houston, W.W.K. & J.A.L. Watson, 1988. Odonata. In: Houston, W.W.K. (ed.) "Zoological catalogue of Australia, volume 6", Australian Government Publishing Service, Canberra, pp. 33-132.
- IUCN, 2003. 2003 IUCN Red List of threatened species. <www.redlist.org/>.
- Lohmann, H., 1996. Das phylogenetische System der Anisoptera (Odonata) [including: Erster Nachtrag]. Entomologische Zeitung 106: 209-252, 253-266, 360-367.
- Malipatil, M.B. & J.D. Blyth, 1982. A qualitative study of the macroinvertebrate fauna of the Thomson River and its major tributaries, Gippsland, Victoria. Reports of the National Museum of Victoria 1: 1-95.
- Marchant, R., 1982. The macroinvertebrates of Magela Creek, Northern Territory. Australian Government Publishing Service, Sydney.
- Marchant, R., P. Mitchell & R. Norris, 1984a. Distribution of benthic invertebrates along a disturbed section of the LaTrobe River, Victoria: an analysis based on numerical classification. Australian Journal of Marine and Freshwater Research 35: 355-374.
- Marchant, R., P. Mitchell & R. Norris, 1984b. A distribution list for the aquatic invertebrates in the lowland region of the LaTrobe River, Victoria. Occasional Papers of the Museum of Victoria 1: 63-79.

- Metzeling, L., A. Graesser, P. Suter & R. Marchant, 1984. The distribution of aquatic macro-invertebrates in the upper catchment of the LaTrobe River, Victoria. Occasional Papers of the Museum of Victoria 1: 1-62.
- Moore, N.W., 1997. Dragonflies – status survey and conservation action plan. IUCN/SSC Odonata Specialist Group. IUCN, Gland & Cambridge.
- New, T.R., 1993. *Hemiphlebia mirabilis* Selys: recovery from habitat destruction at Wilsons Promontory, Victoria, Australia, and implications for conservation management. (Zygoptera: Hemiphlebiidae). Odonotologica 22: 495-502.
- NSW Fisheries, 2002. Adams emerald dragonfly, Fishnote Threatened species in NSW. NSW Fisheries, Nelson Bay.
- O'Connor, N.A., 1993. The fauna of the Pranjip-Creightons Creek system of northern Victoria. Occasional Papers of the Museum of Victoria 6: 54-60.
- Reeves, D.M., 1987. Some dragonflies from Jardine River, Cape York. The Queensland Naturalist 28: 31-32.
- Reeves, D.M., 1988a. Butterflies and dragonflies from Carlisle Island, Cumberland Group. The Queensland Naturalist 29: 21-24.
- Reeves, D.M., 1988b. Dragonflies (Odonata). In: Scott, G. (ed.) "Lake Broadwater – the natural history of an inland lake and its environs", Darling Downs Institute Press, Toowoomba, pp. 170-179.
- Reeves, D.M., 1990. Dragonflies and damselflies (Odonata) from Moreton Island, Queensland. The Queensland Naturalist 30: 71-74.
- Reeves, D.M., 1993. Odonata (dragonflies) from Cape York Peninsula. In: "Cape York Peninsula Scientific Expedition Wet Season Report 1992", The Royal Geographic Society of Queensland Inc., Brisbane, pp. 205-209.
- Reeves, D.M., 1995. Dragonflies and damselflies. In: Ryan, M. (ed.) "Wildlife of Greater Brisbane", Queensland Museum, Brisbane, pp. 55-64.
- Reeves, D.M., 1998. Odonata from Muswellbrook Reserve and Lawn Hill National Park. In: "Muswellbrook Reserve scientific study report 1995", Royal Geographical Society of Queensland Inc., Brisbane, pp. 201-205.
- Reeves, D.M., 2003. Odonata (damselflies and dragonflies) of White Mountains National Park, 8-19 April, 2000. In: "White Mountains scientific study report, 2000", Royal Geographical Society of Queensland Inc., Brisbane, 81-84.
- Reeves, D.M. & P.F. Woodall, 1991. Some Odonata from Eurimbula National Park. The Queensland Naturalist 30: 5-6.
- Sant, G.J. & T.R. New, 1988. The biology and conservation of *Hemiphlebia mirabilis* Selys (Odonata: Hemiphlebiidae) in southern Victoria. Arthur Rylah Institute for Environmental Research Technical Report 82: 1-35.
- Smith, B.J., H.E. Malcolm & P.B. Morison, 1978. Report on the survey of the invertebrate fauna of the Mitta Mitta Valley. In: "Dartmouth Dam Project – report on environmental studies", State Rivers and Water Supply Commission of Victoria: for the River Murray Commission Major Works Branch, Melbourne, pp. 1-90.
- Theischinger, G., 1973. Eine zweite Art der Gattung *Austrocordulia* Tillyard (Odonata: Anisoptera). Annalen des Naturhistorischen Museums Wien 77: 387-397.
- Theischinger, G., 1993. Two new species of *Austroaeschna* Selys from Queensland, Australia (Odonata: Aeshnidae: Brachytroniae). Linzer Biologische Beiträge 25: 805-819.
- Theischinger, G., 1995a. A second species of *Austropetalia* Tillyard from Australia (Odonata: Austropetaliidae). Linzer Biologische Beiträge 27: 291-295.

- Theischinger, G., 1995b. The *Eusynthemis guttata* (Selys) group of species from Australia (Odonata: Synthemistidae). Linzer Biologische Beiträge 27: 297-310.
- Theischinger, G., 1996a. The species of *Austrophlebia* Tillyard (Insecta: Odonata: Anisoptera: Aeshnidae: Brachytroninae). Linzer Biologische Beiträge 28: 305-314.
- Theischinger, G., 1996b. The species of Lestoideinae (Insecta: Odonata: Zygoptera: Lestoideidae). Linzer Biologische Beiträge 28: 315-324.
- Theischinger, G., 1997a. The *Pseudagrion ignifer* complex from Australia (Odonata: Zygoptera: Coenagrionidae). Linzer Biologische Beiträge 29: 799-805.
- Theischinger, G., 1997b. A new species of *Austrosticta* Tillyard from Australia (Insecta: Odonata: Zygoptera: Isostictidae). Linzer Biologische Beiträge 29: 807-810.
- Theischinger, G., 1997c. Two cases of interspecific mating of Australian dragonflies (Anisoptera: Aeshnidae and Corduliidae). Notulae Odonatologicae 4: 164.
- Theischinger, G., 1998a. Supra-specific diversity in Australian "*Argiolestes*" (Odonata: Zygoptera: Megapodagrionidae). Stapfia 55: 613-621.
- Theischinger, G., 1998b. *Tonyosynthemis*, a new dragonfly genus from Australia (Insecta: Odonata: Synthemistidae). Linzer Biologische Beiträge 30: 139-142.
- Theischinger, G., 1998c. A new species of *Griseargiolestes* Theischinger from Australia (Odonata: Zygoptera: Megapodagrionidae). Stapfia 55: 623-627.
- Theischinger, G., 1998d. The *Eusynthemis guttata* (Selys) group of species from Australia (Odonata, Synthemistidae). Linzer Biologische Beiträge 30: 147-153.
- Theischinger, G., 1998e. The larvae of the Australian Gomphidae (Anisoptera). Odonatologica 27: 435-467.
- Theischinger, G., 1998f. A new species of *Eusynthemis* Förster from Australia (Odonata: Synthemistidae). Linzer Biologische Beiträge 30: 143-146.
- Theischinger, G., 1999a. A new species of *Petalura* Leach from south-eastern Queensland (Odonata: Petaluridae). Linzer Biologische Beiträge 31: 159-166.
- Theischinger, G., 1999b. A new gomphid species from the Kimberleys in north-western Australia (Insecta: Odonata). Linzer Biologische Beiträge 31: 369-372.
- Theischinger, G., 1999c. New and little-known Synthemistidae from Australia (Insecta: Odonata). Linzer Biologische Beiträge 31: 373-379.
- Theischinger, G., 1999d. Regions of taxonomic disjunction in Australian Odonata and other freshwater insects: first addendum, with the description of *Austrocordulia refracta juritzai* ssp. nov. (Anisoptera: Corduliidae). Odonatologica 28: 377-384.
- Theischinger, G., 2000a. Preliminary keys for the identification of larvae of the Australian Gomphides (Odonata). Cooperative Research Centre for Freshwater Ecology, Thurgoona.
- Theischinger, G., 2000b. The *Acanthaeschna* story. Linzer Biologische Beiträge 32: 235-240.
- Theischinger, G., 2000c. A new species of *Pseudagrion* Selys from Australia (Odonata: Coenagrionidae). Linzer Biologische Beiträge 32: 253-257.
- Theischinger, G., 2000d. The male of *Eusynthemis tenera* Theischinger (Odonata: Synthemistidae). Linzer Biologische Beiträge 32: 241-245.
- Theischinger, G., 2001a. Preliminary keys for the identification of larvae of the Australian Synthemistidae, Gomphomacromiidae, Pseudocorduliidae, Macromiidae and Austrocorduliidae (Odonata). Cooperative Research Centre for Freshwater Ecology, Thurgoona.
- Theischinger, G., 2001b. Regions of taxonomic disjunction in Australian Odonata and other freshwater insects: second addendum, with the description of *Austroaeschna unicornis pinheyi* ssp. nov. (Anisoptera: Aeshnidae). Odonatologica 30: 87-96.

- Theischinger, G., 2001c. A new species of *Eurysticta* Watson from Australia (Odonata: Isostictidae). Linzer Biologische Beiträge 33: 1291-1294.
- Theischinger, G., 2002. Preliminary keys for the identification of larvae of the Australian Petaluridae, Archipetaliidae, Austropetaliidae, Telephlebiidae and Aeshnidae (Odonata). Cooperative Research Centre for Freshwater Ecology, Thurgooda.
- Theischinger, G., 2003. A new species of *Nannophya* Rambur from Australia (Odonata: Libellulidae). Linzer Biologische Beiträge 35: 661-666.
- Theischinger, G., 2004. Affinities and status of some genus group taxa in Australian Gomphidae (Anisoptera). Odonatologica 33: in press.
- Theischinger, G. & J.H. Hawking, 2003. Dragonflies of Victoria. An identification guide to adult and larval dragonflies (Odonata). Cooperative Research Centre for Freshwater Ecology, Thurgooda.
- Theischinger, G. & J.A.L. Watson, 1978. The Australian Gomphomacromiinae (Odonata: Corduliidae). Australian Journal of Zoology 26: 399-431.
- Theischinger, G. & J.A.L. Watson, 1982. Larvae of the Australian Gomphomacromiinae, and their bearing on the status of the *Synthemis* group of genera (Odonata: Corduliidae). Australian Journal of Zoology 32: 67-95.
- Theischinger, G., J.A.L. Watson & R. Rowe, 1993. Larvae of Australian Synlestidae (Odonata: Zygoptera). Journal of the Australian Entomological Society 32: 113-119.
- Thompson, D.J., 1989. Dragonflies in caves. Cave Science 16: 75.
- Thompson, D.J., 1991. Dragonflies from the western Kimberley region. Western Australian Naturalist 18: 7.
- Tillyard, R.J., 1909a. Studies in life-histories of Australian Odonata. No. 1. The life history of *Petalura gigantea* Leach. Proceedings of the Linnean Society of New South Wales 34: 256-267.
- Tillyard, R.J., 1909b. Studies in life-histories of Australian Odonata. No. 2. Life history of *Diphlebia lestoides* Selys. Proceedings of the Linnean Society of New South Wales 34: 370-383.
- Tillyard, R.J., 1910a. Studies in life-histories of Australian Odonata. No. 3. Notes on a new species of *Phyllopetalia*, with description of nymph and imago. Proceedings of the Linnean Society of New South Wales 34: 697-708.
- Tillyard, R.J., 1910b. Monograph of the genus *Synthemis* (Neuroptera: Odonata). Proceedings of the Linnean Society of New South Wales 35: 312-377.
- Tillyard, R.J., 1910c. On some experiments with dragonfly larvae. Proceedings of the Linnean Society of New South Wales 35: 665-666.
- Tillyard, R.J., 1911a. Studies in life-histories of Australian Odonata. No. 4. Further notes on the life-history of *Petalura gigantea* Leach. Proceedings of the Linnean Society of New South Wales 36: 86-96.
- Tillyard, R.J., 1911b. On the genus *Cordulephya*. Proceedings of the Linnean Society of New South Wales 36: 388-422.
- Tillyard, R.J., 1912. On the genus *Diphlebia*, with descriptions of new species, and life-histories. Proceedings of the Linnean Society of New South Wales 36: 584-604.
- Tillyard, R.J., 1913. Description and life-history of a new species of *Nannophlebia*. Proceedings of the Linnean Society of New South Wales 37: 572-584.
- Tillyard, R.J., 1916. Life-histories and descriptions of Australian Aeshninae: with a description of a new form of *Telephlebia* by Herbert Campion. Journal of the Linnean Society (Zoology) 33: 1-83.

- Tillyard, R.J., 1928. The larva of *Hemiphlebia mirabilis* Selys (Odonata). Proceedings of the Linnean Society of New South Wales 53: 193-206.
- Trueman, J.W.H., G.A. Hoyer, J.H. Hawking, J.A.L. Watson & T.R. New, 1992. *Hemiphlebia mirabilis* Selys: new localities in Australia and perspectives on conservation (Zygoptera: Hemiphlebiidae). Odonatologica 21: 367-374.
- Watson, J.A.L., 1958. A new species of *Petalura* Leach (Odonata) from Western Australia. Proceedings of the Royal Entomological Society London (B) 27: 116-120.
- Watson, J.A.L., 1962. The dragonflies (Odonata) of south-western Australia. A guide to the identification, ecology, distribution and affinities of larvae and adults. Handbook No. 7. Western Australian Naturalist's Club, Perth.
- Watson, J.A.L., 1963. Life history, distribution and ecology in Odonata of south-western Australia. Proceedings of the North Central Branch of the American Association of Economic Entomologists. 18: 130-133.
- Watson, J.A.L., 1967. The larva of *Synthemis leachi* Selys, with a key to the larvae of Western Australian Synthemidae (Odonata). Western Australian Naturalist 10: 86-91.
- Watson, J.A.L., 1968. Australian dragonflies. Australian Natural History 16: 33-38.
- Watson, J.A.L., 1969. Taxonomy, ecology and zoogeography of dragonflies (Odonata) from the north-west of Western Australia. Australian Journal of Zoology 17: 65-112.
- Watson, J.A.L., 1981. Odonata (dragonflies and damselflies). In: Keast, A. (ed.) "Ecological biogeography of Australia, vol. 2", Junk, The Hague, pp. 1139-1167.
- Watson, J.A.L., 1991. The Australian Gomphidae (Odonata). Invertebrate Taxonomy 5: 289-441.
- Watson, J.A.L., 1995. The conservation status of the enigmatic Australian dragonfly *Hemiphlebia mirabilis* Selys. In: Corbet, P.S., S.W. Dunkle, & H. Ubukata (eds) "Proceedings of the International Symposium on the conservation of dragonflies and their habitats [Kushiro, 1993]", Japanese Society for Preservation of Birds, Kushiro, pp. viii, 16-18.
- Watson, J.A.L. & H.M. Abbey, 1980. Dragonflies (Odonata) from the Northern Territory. CSIRO Division of Entomology Report 13: 1-44.
- Watson, J.A.L. & W.W.K. Houston, 1994. Checklist and primary literature for Australian dragonflies (Odonata). Odonatologica 23: 23-44.
- Watson, J.A.L. & G. Theischinger, 1980. The larva of *Antipodophlebia asthenes* (Tillyard): a terrestrial dragonfly? (Odonata: Aeshnidae). Odonatologica 9: 253-258.
- Watson, J.A.L. & G. Theischinger, 1984. Regions of taxonomic disjunction in Australian Odonata and other freshwater insects. Odonatologica 13: 147-157.
- Watson, J.A.L., G. Theischinger & H.M. Abbey, 1991. The Australian dragonflies. A guide to the identification, distributions and habitats of Australian Odonata. CSIRO Australia, Canberra & Melbourne.
- Wells, A. & D.I. Cartwright, 1992. Trichoptera, Ephemeroptera, Plecoptera and Odonata of the Jardine River area, Cape York Peninsula, Northern Queensland. In "Cape York Peninsula Scientific Expedition Report. Volume 2", The Royal Geographical Society of Queensland Inc., Brisbane, pp. 221-230.
- Woodall, P.F., 1992. Some Odonata from south-western Queensland, with notes on relative abundance of Zygoptera at two localities. The Queensland Naturalist 31: 123-125.
- Woodall, P.F., 1993. Odonata from the central Queensland coast: Deepwater and Kinkuna National Parks. The Queensland Naturalist 32: 58-59.